



# Treatment strategies for autoimmune encephalitis

Yong-Won Shin, Soon-Tae Lee, Kyung-Il Park, Keun-Hwa Jung, Ki-Young Jung, Sang Kun Lee and Kon Chu

*Ther Adv Neurol Disord*

2018, Vol. 11: 1–19

DOI: 10.1177/  
1756285617722347

© The Author(s), 2018.  
Reprints and permissions:  
[http://www.sagepub.co.uk/  
journalsPermissions.nav](http://www.sagepub.co.uk/journalsPermissions.nav)

**Abstract:** Autoimmune encephalitis is one of the most rapidly growing research topics in neurology. Along with discoveries of novel antibodies associated with the disease, clinical experience and outcomes with diverse immunotherapeutic agents in the treatment of autoimmune encephalitis are accumulating. Retrospective observations indicate that early aggressive treatment is associated with better functional outcomes and fewer relapses. Immune response to first-line immunotherapeutic agents (corticosteroids, intravenous immunoglobulin, plasma exchange, and immunoabsorption) is fair, but approximately half or more of patients are administered second-line immunotherapy (rituximab and cyclophosphamide). A small but significant proportion of patients are refractory to all first- and second-line therapies and require further treatment. Although several investigations have shown promising alternatives, the low absolute number of patients involved necessitates more evidence to establish further treatment strategies. In this review, the agents used for first- and second-line immunotherapy are discussed and recent attempts at finding new treatment options are introduced.

**Keywords:** autoimmune encephalitis, immunotherapeutic agent, treatment option

Received: 15 March 2017; revised manuscript accepted: 14 June 2017.

## Introduction

Limbic encephalitis (LE), first described in 1960,<sup>1</sup> is characterized by a subacute onset of episodic memory loss and confusion frequently accompanied by seizures, psychiatric symptoms, and lesion involving the medial temporal lobe and hippocampus. Infectious agents such as herpes simplex virus cause inflammation in the central nervous system (CNS) including the limbic area of the brain, but a substantial number of patients with LE are without clear evidence of CNS infection. The existence of autoantibodies and a response to immunotherapy in these patients suggests that an aberrant immune reaction is involved in the disease pathogenesis. Autoimmune etiology is increasingly recognized as a major cause of LE along with the finding of the high prevalence of anti-*N*-methyl-D-aspartate receptor (NMDAR) antibody-associated encephalitis after the discovery of the antibody<sup>2</sup> and the continued identification of additional novel antibodies in LE. These

autoantibodies often affect not only limbic structures but also other cerebral regions, and the term autoimmune encephalitis (AE), rather than autoimmune LE, is used with growing frequency to refer to this disease entity.

AE is classified according to the location of the antigen, either intracellular or on the cell surface, because each classification is associated with different clinical features, especially pertaining to cancer association and immune therapy responsiveness. Antibodies targeting nuclear and cytoplasmic proteins (onconeural antibodies) such as Hu, Ma, and Ri usually accompany malignancy,<sup>3,4</sup> and LE coincident with the detection of these antibodies is termed ‘paraneoplastic LE.’ Patients producing these antibodies respond poorly to immunotherapy, but treatment of the cancer often results in neurological improvements.<sup>4–6</sup> The intracellular location of the antigens, and

Correspondence to:  
**Kon Chu**  
Department of Neurology,  
Comprehensive Epilepsy  
Center, Laboratory for  
Neurotherapeutics,  
Biomedical Research  
Institute, Seoul National  
University Hospital, 101,  
Daehangno, Jongno-  
gu, Seoul 110-744,  
South Korea Program  
in Neuroscience, Seoul  
National University  
College of Medicine, Seoul,  
South Korea  
[stemcell.snu@gmail.com](mailto:stemcell.snu@gmail.com)

**Yong-Won Shin**  
Department of Neurology,  
Comprehensive Epilepsy  
Center, Laboratory for  
Neurotherapeutics,  
Biomedical Research  
Institute, Seoul National  
University Hospital, Seoul,  
South Korea Program  
in Neuroscience, Seoul  
National University  
College of Medicine, Seoul,  
South Korea  
Yeongju Health Center,  
Gyeongsangbuk-do, South  
Korea

**Kyung-Il Park**  
Department of Neurology,  
Comprehensive Epilepsy  
Center, Laboratory for  
Neurotherapeutics,  
Biomedical Research  
Institute, Seoul National  
University Hospital, Seoul,  
South Korea Program  
in Neuroscience, Seoul  
National University  
College of Medicine, Seoul,  
South Korea  
Department of Neurology,  
Seoul National University  
Hospital Healthcare  
System Gnam Center,  
Seoul, South Korea



Soon-Tae Lee  
Keun-Hwa Jung  
Ki-Young Jung  
Sang Kun Lee  
Department of Neurology,  
Comprehensive Epilepsy  
Center, Laboratory for  
Neurotherapeutics,  
Biomedical Research  
Institute, Seoul National  
University Hospital, Seoul,  
South Korea Program in  
Neuroscience, Seoul  
National University  
College of Medicine, Seoul,  
South Korea

the evidence for CD8<sup>+</sup> cytotoxic T-cell-mediated neuronal cell damage in encephalitis coincident with these antibodies, raise the possibility that the humoral immune response could be a nonpathogenic ‘epiphenomenon.’<sup>6</sup>

Neuronal cell-surface antibodies (NSAbs) are another group of antibodies detected in AE. NSAbs target an extracellular epitope, and their corresponding antigens are often synaptic receptors or components of synaptic protein complexes. Anti-NMDAR antibodies are most common, followed by antibodies against leucine-rich glioma inactivated-1 (LGI1).<sup>7–10</sup> The contactin-associated protein like 2 (Caspr2),<sup>11</sup>  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPA),<sup>12</sup> gamma-aminobutyric acid (GABA)-A and -B receptors,<sup>13,14</sup> dipeptidyl-peptidase-like protein-6 (DPPX),<sup>15</sup> and glycine receptor (GlyR)<sup>16</sup> antibodies are other examples of recently identified NSAbs. Unlike encephalitis with antibodies to intracellular antigens, the frequency of cancer in cases involving antibodies targeting cell-surface antigens is, to a varying degree dependent upon the particular type of antibody, lower.<sup>8,17</sup> Several sources of evidence support the theory that the pathogenesis of encephalitis is mainly antibody-mediated in AEs associated with this group of antibodies.<sup>18</sup> Patients producing NSAbs generally show a good response to immunotherapy and have a better overall prognosis.

Of note, glutamic acid decarboxylase 65 (GAD65) is an intracellular enzyme that catalyzes the conversion of GABA, however, AE presenting with anti-GAD65 antibodies is often non-paraneoplastic and shows better immune responsiveness than paraneoplastic LE.<sup>19,20</sup> In addition, anti-GAD65 antibody is related to diverse diseases including AE, cerebellar ataxia, stiff-person syndrome (SPS), and type 1 diabetes.<sup>21</sup> SPS is a rare condition characterized by axial and limb muscle stiffness and painful spasms, and is associated with antibodies against GAD65, GlyR, and amphiphysin.<sup>22</sup> Anti-GlyR antibodies, in particular, are preferentially associated with progressive encephalitis with rigidity and myoclonus (PERM), an aggressive form of SPS spectrum disorder with additional symptoms including brainstem dysfunction, hyperekplexia, sensory symptoms, and dysautonomia.<sup>23</sup> Anti-GlyR antibodies are directed against cell-surface antigens and patients with these antibodies are more responsive to immunotherapy,<sup>23</sup> compared with those with anti-GAD65 (against intracellular antigens) or

anti-amphiphysin antibodies (evidence of T-cell-mediated neuronal cell damage).<sup>24</sup>

Early, appropriate, and intense treatment is important for achieving a good outcome in AE. However, treatment is challenging because the rarity of the disease limits clinical experience and the evidence base guiding the intervention. Moreover, multiple lines of treatment are essential in many cases, and rather experimental options are not infrequently required. In light of these issues, we review here the currently used AE treatment modalities and recent advances in immunotherapy. The present review focuses on AE associated with NSAbs and, without special notation, we will use the term ‘AE’ to refer to AE with NSAbs only.

### Pathogenic mechanism and diagnosis of autoimmune encephalitis

Most T cells that recognize self-antigens are eliminated in the thymus and some self-reactive T cells that escape this process are controlled in the periphery by further deletion, induction of anergy, or suppression by regulatory T cells (Tregs).<sup>25</sup> Self-reactive B cells are also subject to the processes of negative selection, such as deletion, receptor editing and induction of anergy, throughout the development in the bone marrow and spleen.<sup>26</sup> Patients may become prone to autoimmune disease when there is a loss of tolerance to self-antigens, and dysregulated activation of T and B cells. In particular, activation of self-reactive B cells and their subsequent proliferation and differentiation into autoantigen reactive memory B cells and autoantibody-secreting plasma cells, play pivotal roles in antibody-mediated autoimmunity, including AE. Patients with AE respond to the treatments that aim to lower the titers of autoantibodies, such as intravenous immunoglobulin (IVIg) and plasma exchange (PLEX), emphasizing the role of autoantibodies in disease pathogenesis. Animal model studies replicating features of AE by passive transfer of cerebrospinal fluid (CSF) or immunoglobulins (Igs) from anti-NMDAR encephalitis patients to mice provided more direct evidence of autoantibody pathogenicity.<sup>27,28</sup>

Pathogenic roles of autoantibodies include receptor crosslinking and internalization, complement activation and direct disruption of the epitope, as demonstrated in AE by several immunopathologic analyses and *in vitro* studies.<sup>18,29</sup> In the case

of anti-NMDAR encephalitis, receptor internalization and reduced NMDAR currents were observed in rodent hippocampal neurons and HEK293 cells expressing GluN1 and GluN2 subunits of the NMDAR treated with patient antibody.<sup>30,31</sup> Other experiments showed that the autoantibodies bind to a region within the amino terminal domain of GluN1 and cause disrupted interaction with ephrin B2 receptor and channel function.<sup>32,33</sup>

Anti-NMDAR antibodies are predominantly of the IgG1 subclass of IgG.<sup>34</sup> However, activation of complement and neuronal injury is not evident in the brain tissues from patients with anti-NMDAR encephalitis or the passive transfer mouse model.<sup>27,34–36</sup> In contrast, antibodies against LGI1 and Caspr2 are predominantly IgG4<sup>37–39</sup> and complement activation and neuronal loss are also reported in AEs with these autoantibodies.<sup>36,40</sup> IgG4 antibodies form heterodimers that recognize and bind to two different antigens (hetero-bispecific), acting as monovalent antibodies and being unable to crosslink and internalize the target antigen.<sup>41</sup> In addition, IgG4 antibodies have low affinity to Fc $\gamma$  receptor (Fc $\gamma$ R) and are ineffective in activating complement. As IgG1, but not IgG4, is an effective activator of complement, mechanism of the discrepancy in IgG subclass predominance and complement activation is currently unclear. The IgG1-mediated immune response and direct T-cell-mediated toxicity in the pathogenesis of the AEs need further investigation.

The differences in IgG subclass and complement activation have several clinical implications. Patients with anti-LGI1 encephalitis or anti-Caspr2 encephalitis generally show a less severe clinical manifestation and faster recovery than patients with anti-NMDAR encephalitis, which may reflect the limited ability of IgG4. Long-term cognitive impairment and hippocampal atrophy shown in follow-up MRI<sup>38,42–45</sup> indicate irreversible neuronal damage that may be partly attributable to complement activation and cytotoxic T-cell-mediated neuronal injury. Conversely, reversibility of cerebral atrophy in anti-NMDAR encephalitis is consistent with immunopathological findings including the absence of complement and rare T-cell infiltrates.<sup>46,47</sup> With regard to therapy, immunotherapeutic agents targeting complement, such as eculizumab,<sup>48</sup> could be a viable option for AEs with evidence of complement-mediated neuronal toxicity, but not for AEs without.

AE is diagnosed based on clinical characteristics, magnetic resonance imaging, electroencephalography, functional neuroimaging, work-up for systemic tumors, and detection of autoantibodies.<sup>49</sup> Detection of anti-neuronal autoantibodies in the serum/CSF is used to confirm AE and specify the associated antibody. Immunohistochemistry (IHC) on cryopreserved rodent brain sections is widely used as a primary screening method to detect autoantibodies. This method has the advantage in detecting antibodies against a wide range of antigens, and has the potential to detect novel neuronal antigens; however, it lacks the ability to identify specific antibody targets. Cell-based assays (CBAs) are now widely used to detect specific NSAbs. CBA is an immunoassay using cultured cells (i.e. HEK 293) presenting a specific epitope transfected with the complementary DNA, which has the advantage of preserving the conformational epitope.<sup>50</sup> This method led to the discovery of anti-NMDAR antibodies,<sup>51,52</sup> and the subsequent identification of several other NSAbs.<sup>11,12,14,53</sup> Whereas there is recent evidence against the use of radioimmunoassay results for detecting antibodies to the voltage-gated potassium channel complex without confirmation of antibodies to LGI1 and Caspr2,<sup>54,55</sup> NSAb detection using CBAs still shows great usefulness with high sensitivity and specificity, especially for CSF samples.<sup>50</sup> To achieve the highest sensitivity and specificity, testing both serum and CSF using a combination of IHC and CBAs is recommended.<sup>50,56</sup> However, equipment for identifying these antibodies is not widely available in institutions, and it may take several weeks to get the results. Recently proposed clinical diagnostic criteria for AE without antibody testing help to attain early diagnosis in institutions where antibody testing is not readily accessible.<sup>17</sup>

### Treatment overview

Treatment options for AE range from broadly immune-suppressing agents to those targeting processes in antibody-mediated disease pathogenesis. As in most other inflammatory disorders, corticosteroids are used in the treatment of AE, acting to broadly inhibit the inflammatory process. However, corticosteroids possess less specificity for the antibody-mediated immune process, and their efficacy is limited in cases of AE, and they are associated with several systemic side effects. Other lines of treatment address various specific steps in AE's pathogenesis. Therapeutic targets for these treatments include autoantibodies and other

**Table 1.** Therapeutic agents used in autoimmune encephalitis.

Treatment	Regimen
<b>First-line immunotherapy</b>	
Methylprednisolone	1 g daily, for 3–5 days
Intravenous immunoglobulin	2 g/kg over 5 days (400 mg/kg/day)
Plasma exchange/immunoabsorption	1 session every other day for 5–7 cycles
<b>Second-line immunotherapy</b>	
Rituximab	375 mg/m <sup>2</sup> weekly IV infusion for 4 weeks
Cyclophosphamide	750 mg/m <sup>2</sup> monthly for 3–6 months
<b>Alternative therapy</b>	
Tocilizumab	Initially 4 mg/kg, followed by an increase to 8 mg/kg monthly based on clinical response
Low-dose interleukin-2 (aldesleukin)	1.5 million IU/day, 4 subcutaneous injections with 3-week interval
<b>Steroid-sparing agents used for maintenance therapy</b>	
Azathioprine	Initially 1–1.5 mg/kg once daily or divided twice daily, target 2–3 mg/kg/d
Mycophenolate mofetil	Initially 500 mg twice daily, target 1000 mg twice daily
IV, intravenous.	

immune mediators (IVIg and PLEX), B cells and short-lived plasma cells (rituximab), and specific cytokines associated in the autoimmune and inflammatory process [tocilizumab and low-dose interleukin (IL)-2]. Antiproliferative agents targeting lymphocyte proliferation (cyclophosphamide, azathioprine, mycophenolate mofetil, etc.) are also used in refractory cases or to maintain remission.

Despite the numerous treatment options available, there are issues complicating the treatment of AE. First, peripherally activated B cells can cross the blood–brain barrier (BBB) and undergo clonal expansion and differentiation into antibody-secreting plasmablasts and plasma cells.<sup>8,35</sup> Histopathological study and CSF analysis indicate intrathecal production of antibodies in anti-NMDAR encephalitis.<sup>8,34,35,51,57,58</sup> Immunotherapeutic agents such as rituximab cannot permeate the BBB,<sup>59</sup> which may limit the therapeutic effects of these agents. Second, there are long-lived plasma cells (half-life of >6 months) refractory to conventional immunosuppressive agents and agents targeting B or T cells, leading to the chronicity of autoimmunity with persistent autoantibodies.<sup>60</sup> Long-lived

plasma cells may reside in the bone marrow and intrathecal compartment and contribute to refractoriness and relapses. Proteasome inhibitors target these cells, but clinicians have very limited experience with their use in AE.<sup>61–63</sup> Immunotherapeutic agents used in the treatment of AE are listed in Table 1. A summary of major published series reporting the largest number of patients in each NSAbs are presented in Table 2.

AE is often monophasic, and instances of spontaneous recovery without immunotherapy or tumor resection have been reported.<sup>52,64,65</sup> However, recovery from AE is not without sequelae, and AE-related deaths during the acute stage or follow up after discharge have also been noted.<sup>14,38,43,52,66,67</sup> Even if patients survive without immunotherapy, they may suffer a slower recovery requiring prolonged hospitalization. Persistent cognitive impairment observed in long-term follow up suggests irreversible neuronal death and advocate prompt interruption of disease activity.<sup>38,43,44,68,69</sup>

Patients with AE present with variable clinical manifestations, severity, comorbidity status, and

**Table 2.** Summary of representative<sup>a</sup> case series of each NSAb-associated disorder.

Study	Titulaer and colleagues <sup>72</sup>	Arino and colleagues <sup>38</sup>	van Sonderen and colleagues <sup>39</sup>	Hoffberger and colleagues <sup>67</sup>	Pettingill and colleagues <sup>153</sup>	Hoffberger and colleagues <sup>154</sup>	Carvajal-Gonzalez and colleagues <sup>23</sup>	Tobin and colleagues <sup>155</sup>
Associated NSAb	NMDAR	LG11	Caspr2	AMPA <sup>R</sup>	GABA <sub>A</sub> R	GABA <sub>B</sub> R	GlyR	DPPX
No. of patients	577	76	38	22	45	20	45 <sup>b</sup>	20
No. of patients with data available for treatment and outcome	501	48	30 for immunotherapeutic agent, 28 for relapse, and 33 for outcome	21	15 (12 for outcome)	20 (19 for immunotherapeutic agent)	45 <sup>c</sup>	20
Clinical manifestation	577/577: anti-NMDAR encephalitis	63/76: LE 3/76: non-LE <sup>d</sup> 10/76: encephalopathy <sup>e</sup>	16/38: LE 11/38: Morvan syndrome 5/38: Peripheral nerve hyperexcitability 3/38: Cerebellar syndrome 3/38: others <sup>f</sup>	12/22: LE 8/22: limbic dysfunction with multifocal/diffuse encephalopathy 1/22: LE preceded by motor deficits 1/22: psychosis with bipolar features	15/15: variable (including seizures [n = 7], memory impairment [n = 7], confusion or disorientation [n = 4], psychiatric features [n = 5], hallucination [n = 2], anxiety [n = 4])	17/20: LE 1/20: ataxia 1/20: status epilepticus 1/20: opsoclonus-myoclonus syndrome	33/45: PERM 5/45: LE or epileptic encephalopathy 2/45: SPS 2/45: brainstem features 2/45: demyelinating optic neuropathies 1/45: unclear diagnosis	20/20: encephalopathy (with cortical, cerebellar or brainstem manifestations), myelopathy, and autonomic dysfunction
Female	468 (81%)	26 (34%)	4 (11%)	14 (64%)	23 (51%)	8 (40%)	21 (47%)	8 (40%)
Age, year, median (range)	21 (1–85)	61 (32–80)	66 (25–77)	62 (23–81)	51 (2–80)	61.5 (16–77)	50 (1–75)	53 (13–75)
Tumor	220/577 (38%)	5/76 (7%)	7/37 (19%)	14/22 (64%)	3/14 (21%)	10/20 (50%)	9/45 (20%)	2/20 (10%)
Tumor type	Ovarian teratoma (n = 207), extraovarian teratoma (n = 4), and other tumors (n = 9) <sup>g</sup>	Prostate cancer (n = 2), gastric neuroendocrine tumor (n = 1), colon carcinoma (n = 1), and bone metastasis with unknown origin (n = 1)	Thymoma (n = 4), adenocarcinoma of the lung (n = 1), carcinoma in situ of sigmoid (n = 1), and thoracic mass without pathologic diagnosis (n = 1)	Small-cell lung cancer (n = 5), thymoma (n = 4), breast cancer (n = 2), ovarian teratoma (n = 2), and lung cancer (n = 1)	Dysembryoplastic neuroepithelial tumor (n = 1), prostatic cancer (n = 1), and Non-Hodgkin's lymphoma (n = 1)	Small-cell lung cancer (n = 10)	Thymoma (n = 3), follicular lymphoma and previous treated cancer (n = 5) <sup>h</sup>	Gastrointestinal follicular lymphoma (n = 1) and chronic lymphocytic leukemia (n = 1)
First-line immunotherapy	462/501 (92%)	48/48 (100%)	28/30 (93%) without tumor <sup>i</sup>	20/21 (95%)	4/15 (27%)	15/19 (79%)	37/44 (84%)	11/20 (55%)
Corticosteroids	421/501 (84%)	44/48 (92%)	16 / 30 (53%)	17/21 (81%)	2/15 (13%)	14/19 (74%)	31/44 (70%)	11/20 (55%)
IVIg	346/501 (69%)	26/48 (54%)	13 / 30 (43%)	12/21 (57%)	1/15 (7%)	7/19 (37%)	20/44 (45%)	5/20 (25%)

(Continued)



**Table 2.** (Continued)

Study	Titulaer and colleagues <sup>72</sup>	Arino and colleagues <sup>38</sup>	van Sonderen and colleagues <sup>39</sup>	Hoffberger and colleagues <sup>67</sup>	Pettingill and colleagues <sup>153</sup>	Hoffberger and colleagues <sup>154</sup>	Carvajal-Gonzalez and colleagues <sup>23</sup>	Tobin and colleagues <sup>155</sup>
PLEX	163/501 (33%)	0/48 (0%)	3/30 (10%)	6/21 (29%)	2/15 (13%)	5/19 (26%)	17/44 (39%)	5/20 (25%)
Second-line immunotherapy	134/501 (27%)	7/48 (15%)	7/30 (23%)	5/21 (24%)	0/15 (0%)	3/19 (16%)	4/44 (9%)	7/20 (35%)
Rituximab	101/501 (20%)	6/48 (13%)	5/30 (17%)	5/21 (24%)	0/15 (0%)	2/19 (11%)	2/44 (5%)	5/20 (25%)
Cyclophosphamide	81/501 (16%)	3/48 (6%)	2/30 (7%)	1/21 (5%)	0/15 (0%)	1/19 (5%)	4/44 (9%)	3/20 (15%)
Other immunotherapy	31/501 (6%): azathioprine, mycophenolate mofetil, tacrolimus or methotrexate	7/48 (15%): azathioprine (n = 6), mycophenolate mofetil (n = 1)	1/30 (3%): azathioprine	0/21 (0%)	1/15 (7%): azathioprine	1/19 (5%): mycophenolate mofetil	7/44 (16%): azathioprine (n = 4), mycophenolate mofetil (n = 3), cyclosporine (n = 1)	3/20 (15%): azathioprine (n = 2), mycophenolate mofetil (n = 1)
Follow up, median (range)	24 months (4–186)	39 months (22–58)	36 months (range 3–168)	72 weeks (5–266)	18 months (2–20)	7 months (0.75–45)	3 years (18 months to 7 years)	6 months (0–68)
Relapse	45/501 (9%)	10/48 (21%)	7/28 (25%) <sup>i</sup>	1/21 (5%)	1/12 (8%)	Not provided	6/43 (14%)	2/20 (10%)
Outcome	mRS 0–2: 394/501 (79%) mRS 3–5: 77/501 (15%) mRS 6: 30/501 (6%)	CPS0-1: 34 (71%) CPS2-4: 14 (29%)	mRS 0-2: 24/33 (73%) mRS 3-5: 5/33 (15%) mRS6: 4/33 (13%)	mRS 0-2: 10/21 (48%) mRS 3-5: 6/21 (29%) mRS 6: 5/21 (24%)	Improved: 8/12 (67%) Declined: 1/12 (8%) Huntington disease confirmed: 1/12 (8%) Death: 2/12 (17%)	Treatment response - Complete response: 7/16 (44%) - Partial response: 8/16 (50%) - No response: 1/16 (6%) Death: 8/20 (40%)	mRS 0-2: 34/44 (77%) mRS 3-5: 6/44 (14%) mRS 6: 4/44 (9%)	Complete remission or mild disability: 4/18 (22%) Partial response to treatment: 5/18 (28%) Unchanged: 6/18 (33%) Worsened: 3/18 (6%) including 2 deaths

<sup>a</sup>Largest in number of patients and fully dedicated to a single NSAb.

<sup>b</sup>A total of seven patients with very low GlyR antibodies (not titrating beyond 1:40) were excluded.

<sup>c</sup>Data on each of immunotherapeutic agent, relapse, and outcome are not available for 1 or 2 patients.

<sup>d</sup>There were two that had MRI abnormality outside temporal lobes, one with normal MRI with CSF pleocytosis.

<sup>e</sup>No MRI or CSF evidence of inflammation.

<sup>f</sup>Single seizure followed by pain syndrome (n = 1), painful polyneuropathy (n = 1), or mild amnesic syndrome with frontal lobe dysfunction (n = 1).

<sup>g</sup>Lung (n = 2), breast (n = 2), testicular (n = 2), ovarian (n = 1), thymic (n = 1), and pancreatic (n = 1) carcinoma.

<sup>h</sup>Breast cancer (n = 1), breast cancer and thymoma (n = 1), thymoma and lymphoma (n = 1), Hodgkin lymphoma (n = 1), malignant melanoma (n = 1).

<sup>i</sup>Overall, four of seven patients with tumor were treated with immunotherapy, but information for immunotherapeutic agent is not provided.

<sup>j</sup>Patients with a ≥ 1 year follow up.

AMPAR, α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor; Caspr2, contactin-associated protein like 2; CPS, Cognitive Performance Scale; CSF, cerebrospinal fluid; DPPX, dipeptidyl-peptidase-like protein-6; GABA<sub>A</sub>R, gamma-aminobutyric acid A receptor; GABA<sub>B</sub>R, gamma-aminobutyric acid B receptor; GlyR, glycine receptor; IVIg, intravenous immunoglobulin; LE, limbic encephalitis; LGI1, leucine-rich glioma inactivated-1; MRI, magnetic resonance imaging; mRS, modified Rankin Scale; NMDAR, N-methyl-D-aspartate receptor; NSAb, neuronal cell-surface antibody; PERM, progressive encephalomyelitis with rigidity and myoclonus; PLEX, plasma exchange; SPS, stiff-person syndrome.

immunotherapy responsiveness, and thus treatment should be individualized. There are no established guidelines for treatment, and diverse regimens are currently being used based on the patient's clinical status and the clinician's opinion. The three common factors derived from a recent systematic review for better outcomes and fewer relapses are the commencement of immunotherapy, early initiation of therapy, and commencement of second-line immunotherapy if first-line immunotherapy fails.<sup>70</sup> Additionally, a thorough tumor screening in the early stages of the disease is important for desirable patient outcomes. Tumors such as ovarian teratoma, thymoma, and small-cell lung carcinoma are detected in a considerable proportion of AE cases.<sup>17</sup> Ovarian teratomas, formerly believed to be benign tumors, are strongly associated with anti-NMDAR encephalitis, and resection of the tumor is an important part of the treatment.<sup>52,71,72</sup>

### First-line immunotherapy

Common first-line immunotherapeutic agents include corticosteroids, IVIg, and PLEX. Although there is no compelling evidence to suggest the superiority of any specific regimen, corticosteroids are frequently the first choice, followed by IVIg and PLEX.<sup>70</sup> Corticosteroids with either IVIg or PLEX represent the usual choice when a combination of first-line agents is administered. For pediatric patients and patients with medical conditions that discourage the use of corticosteroids, such as diabetes mellitus and seropositivity for type 1 diabetes-associated antibodies such as GAD65,<sup>73</sup> IVIg is often the initial treatment. PLEX is not easily performed on pediatric patients, patients with autonomic instability, or those who cooperate poorly.<sup>66</sup> However, PLEX is useful when patients poorly tolerate corticosteroids and IVIg. The clinical effectiveness of each of these agents is nearly confirmed by previous observations, but more formal studies are necessary to identify the ideal first-line treatment regimen.

### Corticosteroids

Corticosteroids bind to intracellular glucocorticoid receptors and suppress the transcription of multiple proinflammatory genes that encode cytokines, chemokines, adhesion molecules, inflammatory enzymes, receptors and proteins.<sup>74</sup> Glucocorticoids have an influence on almost all cytokines, and their use results in the depletion of

T cells, inhibition of Th1 differentiation, macrophage dysfunction, and eosinophil apoptosis.<sup>75</sup> At higher concentration, corticosteroids have additional effects on the synthesis of anti-inflammatory proteins, and also induce post-transcriptional effects.<sup>74</sup> Furthermore, corticosteroids offer extra benefit to CNS inflammatory disorders by restoring BBB integrity and controlling brain edema.<sup>76</sup>

While it is apparent from observations of the last decade that corticosteroids have therapeutic effects in cases of AE, there are several issues that should be considered in their use. First, the differentiation between AE and infectious encephalitis is often difficult in the acute stage, which frequently delays the initiation of corticosteroid therapy. Second, the reduction of the number of circulating B cells by glucocorticoids is much less than that of T cells,<sup>77</sup> and the effect on serum antibody titer is also limited.<sup>75,78,79</sup> Given the largely antibody-mediated disease pathogenesis,<sup>18</sup> it should be considered that a combination of corticosteroids with other immunotherapeutic agents targeting Igs and B cells may be required for a more effective treatment of AE. Third, along with various medical complications, corticosteroids may induce or aggravate psychiatric symptoms associated with AE, such as depression, insomnia, agitation, and psychosis. Clinical and experimental studies also indicate neurotoxic effects of corticosteroids and their potential to induce neurodegeneration upon chronic exposure.<sup>80</sup> As corticosteroids are the most commonly used agents in first-line immunotherapy for AE, investigations seeking to optimize the use of corticosteroids for better outcomes should be conducted. Clinicians must leverage the clinical benefits and harmful effects of steroids in the AE treatment pathway.

### IVIg

IVIg is a blood product extracted from the collected pool of plasma from over a thousand donors. IVIg provides antibodies to a broad range of pathogens, and is used to provide passive immunity for patients with immunodeficiency.<sup>75</sup> High-dose IVIg (1–2 g/kg) provide various anti-inflammatory and immunomodulatory effects by multidirectional mechanisms such as autoantibody neutralization, blockade of activating FcγR and upregulation of inhibitory FcγRIIB, inhibition of complements, cytokines, and leukocyte migration.<sup>81</sup> In addition, IVIg saturates neonatal Fc receptor (FcRn), a homeostatic regulator of IgG catabolism, by

rescuing IgG from lysosomal degradation, through competition, resulting in the acceleration of IgG breakdown.<sup>82,83</sup> FcRn also contributes to the long half-life of IVIg, and replacement of IVIg is usually considered at 3–4 week intervals. IVIg has been shown to be effective in several autoimmune and inflammatory disorders, and indications for neurological disorders include chronic inflammatory demyelinating polyneuropathy, multifocal motor neuropathy, Guillain-Barré syndrome, and myasthenia gravis.<sup>84</sup>

IVIg can be used as a monotherapy in the treatment of AE, but is more often used after or in combination with high-dose steroids, or with PLEX, rituximab, or other immunotherapeutic agents. IVIg has a better side effect profile than corticosteroids and is more convenient and cost-effective compared with PLEX. However, their comparable efficacy to other first-line agents, and their clinical benefit as an add-on therapy, remains to be clearly established in AE. Considering the wide use of IVIg, and the perceived therapeutic effect according to clinical experiences, controlled studies are needed to establish the utility of IVIg in AE. A multicenter randomized trial for the use of IVIg in pediatric patients with infectious or immune-mediated encephalitis was launched in the UK only recently.<sup>85</sup>

Most of the adverse effects associated with IVIg are mild and transient. There is a risk of anaphylaxis in patients with selective IgA deficiency, but the incidence is extremely rare. Screening of IgA antibody level prior to the administration of IVIg could be helpful but is not considered mandatory.

#### *PLEX and immunoadsorption*

PLEX effectively removes autoantibodies and other pathologic substances in the plasma. PLEX also alters the immune system by changing lymphocyte numbers and their distribution, T-suppressor cell function, and T-helper cell phenotypes.<sup>86</sup> Steroids alone are frequently insufficient to ameliorate autoantibody-mediated immune process, and direct removal or neutralization of autoantibodies from the circulation by PLEX and IVIg may show a synergistic effect. In addition, PLEX increases the proliferation of antibody-producing cells and this could increase susceptibility of these cells to immunosuppressants and chemotherapeutic agents.<sup>86</sup> A small retrospective study in anti-NMDAR encephalitis patients showed that there was greater clinical

improvement with patients who received PLEX immediately after steroids compared with patients who received corticosteroids alone.<sup>87</sup> A recent systematic review by Suppiej and colleagues for PLEX in pediatric anti-NMDAR encephalitis suggested a trend toward better outcomes when PLEX is administered early and when given in combination with steroids.<sup>88</sup>

Immunoadsorption is a refined form of PLEX that enables the selective removal of Igs from separated plasma.<sup>89</sup> Recent studies have shown that immunoadsorption could be an effective therapeutic modality as a part of first-line immunotherapy.<sup>90,91</sup> Equivalent efficacy between immunoadsorption and PLEX was reported in a small study with 21 AE patients with either NSAb or onconeural antibodies.<sup>92</sup>

Technically, PLEX does not remove antibodies from the CSF unless the BBB is severely damaged, and it is unclear whether the removal of systemic antibodies effectively decreases CSF antibody levels. However, recent reports on the use of immunoadsorption for the treatment of AE demonstrated a 64% decrease in CSF antibody titers at early follow up (median 5 days after the last session).<sup>91</sup> Transient elevation of intrathecal IgG fraction and IgG index ('spurious quantitative intrathecal Ig synthesis') is frequently observed in the first 2 days following PLEX or immunoadsorption.<sup>93</sup> However, this does not mean true elevation of intrathecal Ig synthesis, and clinicians should be aware of this phenomenon to avoid unnecessary diagnostic and therapeutic procedures.

#### **Second-line immunotherapy**

First-line immunotherapy is often insufficient in the treatment of AE, and secondary immunomodulatory agents are typically used. Observational studies show that second-line treatment results in better functional outcomes and lower relapse rates with manageable adverse effects.<sup>52,66,72,94</sup> The decision to initiate second-line immunotherapy is made with consideration of the disease severity, response to the first-line immunotherapy, presence of relapse, and other clinical conditions. Consensus criteria on the appropriate time to initiate second-line immunotherapy are yet to be established, but a quick progression to second-line immunotherapy is favored in some institutions including in our group.<sup>94,95</sup> Rituximab and cyclophosphamide are the most



commonly used second-line agents in AE treatment. However, these agents are not without their shortcomings, and are not effective in all patients. Additional therapeutic options should be introduced in the second-line immunotherapy but clinical experience is currently insufficient to permit doing so.

### *Rituximab*

Rituximab is a partially humanized monoclonal antibody directed against CD20, a glycoprotein primarily found on the surface of B cells, initially approved for the treatment of non-Hodgkin B-cell lymphomas. It is widely used to treat various autoimmune disorders and appears to be effective in several autoimmune CNS and peripheral nervous system disorders.<sup>96</sup> Rituximab depletes both naïve and memory B cells through antibody-mediated cellular toxicity, complement activation, and induction of apoptosis.<sup>96</sup> A substantial reduction in relapse rate with the depletion of memory B cells has been demonstrated in neuromyelitis optica (NMO) or NMO spectrum disorder (NMOSD).<sup>97–99</sup> Circulating levels of B cells are usually below the detectable range for 6–8 months after treatment.<sup>100</sup> Effective depletion of B cells in the peripheral blood is confirmed in AE patients,<sup>94,101</sup> and subsequent depletion of short-lived plasmablasts is also reported in anti-NMDAR encephalitis.<sup>102</sup> The high effectiveness of rituximab in IgG4-related disease<sup>103</sup> further supports the use of rituximab in AE in which antibodies of IgG4 subclass predominate, such as in anti-LGI1,<sup>37,38</sup> anti-Caspr2,<sup>37,39</sup> and anti-IgLON5 encephalitis.<sup>104</sup> Rituximab does not target CD20-negative cells including long-lived plasma cells, thus reducing adverse immunosuppressive effects.<sup>100,102</sup>

Lee and colleagues reported on the efficacy and safety of rituximab as a second-line immunotherapy for AE.<sup>94</sup> In this retrospective study of 161 patients, additional rituximab treatment was associated with improvement of functional outcomes measured by the modified Rankin Scale (mRS). This study included AE with or without proven antibody (both NSAb and onconeural antibodies) status, and showed rituximab to be effective independent of patient antibody status. Infusion-related adverse effects were noted in 5 (6.7%) patients and infection (all pneumonia) in 9 (11.3%), but there were no life-threatening or recurrent infectious adverse events. Given the substantial efficacy and safety of rituximab in the

treatment of AE, and the known association of better outcomes with early treatment,<sup>72</sup> the authors proposed the incorporation of rituximab into the first-line treatment protocol. Additionally, one study of the use of rituximab in a pediatric population with autoimmune or inflammatory CNS disease further supported the administration of rituximab in the early stages of the disease.<sup>101</sup> However, this study reported 11 (7.6%) patients with infectious complications, including two life-threatening or disabling infections and two deaths. The authors concluded that rituximab should be reserved only for those with significant morbidity and risk of mortality in the pediatric population.

Rituximab therapy increases the risk of reactivation of chronic viral infections such as hepatitis B, and serologic screening tests should be considered prior to initiation of the treatment.<sup>105</sup> Progressive multifocal leukoencephalopathy is a rare CNS complication of rituximab therapy,<sup>106</sup> and has not yet been reported in patients with AE.

### *Cyclophosphamide*

Cyclophosphamide is an alkylating agent that inhibits cell proliferation, affecting both B and T cells. Along with rituximab, it is a constituent of the so-called R-CHOP regimen (rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisolone) used for the treatment of non-Hodgkin lymphoma.<sup>107</sup> It is a widely used chemotherapeutic agent as well as an immunosuppressant for life-threatening or severe rheumatologic and renal diseases such as ANCA-associated vasculitis, lupus nephritis, and other systemic vasculidites.<sup>75,107–109</sup> However, it is not typically used to treat autoimmune neurological disorders and is a less preferable agent than rituximab as a second-line agent in AE. Its potentially serious side effects such as myelosuppression, infertility, hemorrhagic cystitis, and an increased risk of malignancy lower the priority of its use. However, its low cost (compared with rituximab), direct suppression of lymphocyte proliferation (unlike first-line agents), and the greater accumulated clinical experience of its use (compared with the immunotherapeutic agents currently not included in the first- or the second-line therapies) contribute to the justification of its use in refractory cases. Gonadotropin-releasing hormone agonist administration or egg/sperm collection may be employed to preserve fertility following cyclophosphamide use.<sup>110</sup>

### Alternative therapy

Approximately 20–50% of patients with AE show inadequate responses to second-line therapies, and exhibit persistent neurological issues.<sup>72,94</sup> Re-administration of first-line immunotherapeutic agents, extended use of second-line immunotherapy, and long-term maintenance of prednisolone or steroid-sparing agents such as azathioprine and mycophenolate mofetil, are among the options that have been used so far. Mycophenolate mofetil, in particular, has selective antiproliferative activity on lymphocytes and has shown better efficacy in inducing remission and a more favorable side effect profile than cyclophosphamide in other autoimmune disorders,<sup>111</sup> supporting its use as a safer alternative to cyclophosphamide for second-line immunotherapy in AE. A small number of studies reported a more targeted therapy with monoclonal antibodies or direct infusion of immune mediators.<sup>61–63,112,113</sup>

### Tocilizumab

Tocilizumab is a monoclonal antibody targeting the IL-6 receptor. IL-6 induces B-cell proliferation and differentiation into antibody-producing cells, promotes differentiation of CD8<sup>+</sup> cytotoxic T cells, induces differentiation of naïve CD4<sup>+</sup> helper T cells into IL-17-producing T-helper cells, and inhibits differentiation of those cells into regulatory T cells, which all contribute to autoimmune tissue damage.<sup>114</sup> Tocilizumab binds to both soluble and membrane bound IL-6 receptors and inhibits IL-6 from binding to its receptors, leading to the blockade of IL-6-mediated inflammatory cascades.<sup>115</sup> The therapeutic effect of tocilizumab is demonstrated in various autoimmune diseases, including rheumatoid arthritis<sup>116</sup> and systemic juvenile idiopathic arthritis.<sup>117</sup> Among autoimmune neurological disorders, the efficacy of tocilizumab in therapy-resistant cases of NMO and NMOSD was demonstrated in several small studies.<sup>118–120</sup> A case report of anti-Caspr2 encephalitis successfully treated with tocilizumab showed therapeutic potential of tocilizumab for therapy-resistant AE.<sup>121</sup>

A recent observational study by Lee and colleagues showed that tocilizumab potentially improves clinical symptoms of AE in patients who do not respond adequately to rituximab.<sup>112</sup> In this study, the patients who received tocilizumab showed superior clinical improvement at 1 month

and at the last follow up compared with patients who received additional monthly rituximab or observation without additional immunotherapy. These results encourage the introduction of new agents with different mechanisms of action for use in refractory cases. However, the study was conducted retrospectively without standardized treatment and follow-up protocols, and 65.9% of the patients studied had seronegative AE. These results must therefore be confirmed by further studies with more sophisticated design and larger samples in order to establish tocilizumab as a viable alternative treatment option for therapy-resistant cases of AE.

Tocilizumab increases the risk of infection, but impaired IL-6 receptor signaling may blunt the fever response and elevation of C-reactive protein.<sup>122,123</sup> Clinicians need to be wary of systemic infection in patients treated with tocilizumab. Other side effects include neutropenia, thrombocytopenia, and elevated liver enzymes and lipid levels. Regular checkup of complete blood count with differential, liver profile, and lipid levels is required after tocilizumab treatment.

### Low-dose IL-2 therapy and Treg modulation

The number and function of Tregs are dysregulated in autoimmune conditions.<sup>124,125</sup> IL-2 is a key regulator for Treg differentiation, survival, and function, and has a role in keeping tolerance over autoimmunity.<sup>126,127</sup> IL-2 activates effector and memory T cells, and is classified as a proinflammatory cytokine. However, low-dose IL-2 administration can selectively expand Tregs without promoting effector T-cell responses because Tregs have lower activation thresholds to IL-2 than to effector T cells.<sup>128,129</sup> In this regard, low-dose IL-2 therapy has been proposed as a promising new therapeutic option for autoimmune and inflammatory disorders with Treg insufficiency.<sup>130</sup> Some studies report that low-dose IL-2 administration was clinically beneficial, with improvement of Treg number and functions in hepatitis C virus-induced vasculitis,<sup>131</sup> alopecia areata,<sup>132</sup> systemic lupus erythematosus,<sup>133</sup> and graft-versus-host disease.<sup>134</sup>

Lim and colleagues evaluated the feasibility of low-dose IL-2 therapy with 10 patients with refractory AE.<sup>113</sup> The patients underwent four to five cycles of low-dose IL-2 therapy and showed modest responses with a median mRS improvement from three to two. All four patients with

anti-NMDAR encephalitis responded to the therapy, and two of six antibody-negative AE patients showed seizure reduction. Serious side effects were observed in two patients (decreased neutrophil count and ileus) but these events did not lead to the termination of the treatment. Increased eosinophil counts were common (8 of 10 patients) but resolved spontaneously. These results warrant further trials to clarify the efficacy of low-dose IL-2 therapy in AE.

### *Bortezomib*

Bortezomib is a proteasome inhibitor particularly effective at depleting plasma cells and is approved for the treatment of multiple myeloma.<sup>135</sup> Given that long-lived plasma cells are not the target of B-cell-depleting agents and are also resistant to glucocorticoid and antiproliferative agents such as cyclophosphamide,<sup>60</sup> bortezomib and other proteasome inhibitors can represent alternative options for refractory cases. Currently, three reports with nine cases exist that have explored the treatment of severe anti-NMDAR encephalitis with bortezomib.<sup>61–63</sup> Clinical improvement as well as a tolerable safety profile was reported with bortezomib use in these patients. However, a prolonged treatment period or intensive treatment combined with several immunotherapeutic agents obscured the effectiveness of bortezomib therapy, and the remission achieved could be part of the natural course of the disease or the effect of preceding agents. More experience with bortezomib therapy is needed before making any conclusions.

### **Maintenance therapy**

AE is not always monophasic and relapse of AE has been noted even after 5–10 years.<sup>38,43,136</sup> The rate of relapse stated in recent literature (approximately 10–20%, varying with the type of antibody<sup>70</sup>) may still be underestimated, given the relatively short time period since the discovery of the relevant autoantibodies. Early aggressive therapy is reported to reduce relapse rates,<sup>70,72</sup> but the role of maintenance therapy is largely unexplored. Overlap with multiple sclerosis,<sup>137–139</sup> NMO or NMOSD,<sup>140,141</sup> and myelin oligodendrocyte glycoprotein antibody positive demyelinating syndrome<sup>141</sup> in anti-NMDAR encephalitis suggests that long-term maintenance therapy is required at least for a specific population. Maintenance therapy is usually considered in clinical practice in order to maximize therapeutic gain and attain the

highest functional state possible, as well as to make certain that the patient reaches complete remission free of relapse. One study found that additional monthly rituximab after 4 weekly infusions contributed to further mRS improvement.<sup>94</sup> Sustained use of oral corticosteroids and monthly IVIg or PLEX may also be considered for patients with incomplete recovery. Azathioprine and mycophenolate mofetil are commonly used oral steroid-sparing agents for maintenance therapy in autoimmune neurological disorders such as myasthenia gravis and NMO.<sup>142,143</sup> These agents may be used following acute treatment of AE for sustained remission, however, it remains to be evaluated whether long-term immunosuppression with oral agents is effective in reducing the relapse rate of AE. Intermittent administration of rituximab at regular 6-month intervals or depending on circulating B-cell numbers may be considered as another option for patients who respond well to rituximab therapy. The appropriate duration of maintenance therapy is currently unknown and the length of empirical use ranges widely from 6 months to several years according to the patient's status and clinician's opinion.

### **Future directions**

Criteria and guidelines for the clinical diagnosis of AE have recently been developed by expert consensus.<sup>17</sup> However, there have been no randomized controlled trials for the treatment of AE, and immunotherapeutic agents currently used in AE do not have a definite indication due to the low level of supporting evidence. Given the rarity of AE, international collaboration for prospective clinical trials is imperative to establish treatment guidelines. There are several additional issues that should be considered to establish better treatment strategies. First, an optimized regimen and dosing schedule is not yet determined. Regimens used for acute management differ among clinicians, and there is no clear standard to determine treatment failure and when to initiate different immunotherapeutic agents. The appropriate duration of immunotherapy for sustained remission must also be determined. Second, while treatment response, length of treatment, and outcome vary depending on the type of associated antibody, current immunotherapeutic regimens for AE are not tailored to the type of associated antibody. Third, the proportion of patients who show inadequate response to first- and second-line immunotherapy is not negligible, and yet several new

immunotherapeutic agents are still not applied in the treatment of AE. Investigation of more therapeutic options is needed to augment the therapeutic armamentarium for the treatment of AE. Attempt to induce immune tolerance, for instance, is an option that is yet poorly explored for the treatment of AE. In NMOSD and other autoimmune disorders, several innovative approaches to induce antigen-specific tolerance or other strategies to restore central and peripheral tolerance are being developed.<sup>144,145</sup>

Lastly, more investigations into biomarkers for AE are needed to aid early diagnosis and to predict treatment response, relapse, and outcomes. CSF profile and oligoclonal band positivity differ among associated antibodies.<sup>8,10</sup> There is also a report that a low white blood cell count on the first CSF examination is associated with a good neurologic outcome at 6 months after admission in adult patients admitted to intensive care unit with anti-NMDAR encephalitis.<sup>146</sup> However, the association between CSF profile and clinical outcome was not evident in other studies.<sup>57,147</sup> In anti-NMDAR encephalitis, high antibody titer is associated with poor outcome and/or the presence of teratoma, and faster and greater decrease in CSF antibody titer during the first 1 month of the disease is associated with better outcome.<sup>56</sup> Recovery and relapses are associated with decrease and increase of CSF antibody titer, respectively. However, the correlation between antibody titers and clinical symptoms is imperfect and the titer of autoantibodies often remains detectable long after remission.<sup>38,56,148</sup> The prognostic utility of the persistence or drop in antibody titer for future relapses is uncertain. It should also be noted that antibody titer is not associated with long-term cognitive outcome in anti-LGI1 encephalitis<sup>38</sup> and the association between antibody titer and clinical outcome in AE with other NSAbs are not well-established. C-X-C motif chemokine 13 (CXCL13) is a B-cell-attracting chemokine, and its levels correlate with the presence of CSF B cells, plasmablasts and intrathecal Ig synthesis.<sup>149</sup> In a study of anti-NMDAR encephalitis, the level of CSF CXCL13 was found to be increased during the early stage of the disease and a high level of CXCL13 was associated with limited response to therapy and clinical relapse.<sup>150</sup> However, CXCL13 is not a specific biomarker for anti-NMDAR encephalitis and was shown to be increased in infectious and demyelinating encephalitis.<sup>151</sup> Another study with CSF biomarkers for

neuronal (neurofilament light chain protein and total tau protein) and glial cell (glial fibrillary acidic protein) damage suggested that these markers reflect disease activity and long-term disability in AE.<sup>152</sup> Investigations of additional chemokine/cytokines and other biomarkers for disease activity may provide valuable information to the clinician in the treatment of AE.

## Conclusion

Although AE responds to immunotherapy and the majority of patients recover from the self-destructive autoimmune process, many patients fail to regain baseline cognitive and functional status. Early aggressive therapy is recommended in AE but steroid abuse should be avoided to prevent potential cognitive and other adverse effects. Each immunotherapeutic agent has its own strengths and weaknesses, and an appropriate combination of these agents is often needed to complement each other and achieve synergistic effects. Results from several studies favor the commencement of second-line therapy and demonstrate the therapeutic potential of a few alternative treatment options. However, the number of patients included in such studies is often very small, and the studies are uncontrolled and prone to severity bias due to the predominant use of these agents in more severe cases. Prospectively designed controlled trials are needed to confirm the efficacy and safety of currently proposed immunotherapeutic agents, along with efforts to uncover more therapeutic options and biomarkers.

## Funding

This work was supported by Seoul National University Hospital Research Fund, South Korea (grant number 2520140040). K Chu was supported by research grants from JW Pharmaceutical (grant number 0620162210), Yuhan (grant numbers 0620151690 and 0620140910), and Ildong Pharmaceutical (grant number 0620130950).

## Conflict of interest statement

The authors declare that there is no conflict of interest.

## References

1. Brierley JB, Corsellis JAN, Hierons R, *et al.* Subacute encephalitis of later adult life - mainly



- affecting the limbic areas. *Brain* 1960; 83: 357–370.
2. Gable MS, Sheriff H, Dalmau J, *et al.* The frequency of autoimmune N-methyl-D-aspartate receptor encephalitis surpasses that of individual viral etiologies in young individuals enrolled in the California Encephalitis Project. *Clin Infect Dis* 2012; 54: 899–904.
  3. Corsellis JA, Goldberg GJ and Norton AR. “Limbic encephalitis” and its association with carcinoma. *Brain* 1968; 91: 481–496.
  4. Gultekin SH, Rosenfeld MR, Voltz R, *et al.* Paraneoplastic limbic encephalitis: neurological symptoms, immunological findings and tumour association in 50 patients. *Brain* 2000; 123: 1481–1494.
  5. Dalmau J and Rosenfeld MR. Paraneoplastic syndromes of the CNS. *Lancet Neurology* 2008; 7: 327–340.
  6. Melzer N, Meuth SG and Wiendl H. Paraneoplastic and non-paraneoplastic autoimmunity to neurons in the central nervous system. *J Neurol* 2013; 260: 1215–1233.
  7. Lancaster E, Martinez-Hernandez E and Dalmau J. Encephalitis and antibodies to synaptic and neuronal cell surface proteins. *Neurology* 2011; 77: 179–189.
  8. Ramanathan S, Mohammad SS, Brilot F, *et al.* Autoimmune encephalitis: recent updates and emerging challenges. *J Clin Neurosci* 2014; 21: 722–730.
  9. Kim SY, Choi SA, Ryu HW, *et al.* Screening autoimmune anti-neuronal antibodies in pediatric patients with suspected autoimmune encephalitis. *J Epilepsy Res* 2014; 4: 55–61.
  10. Byun JI, Lee ST, Jung KH, *et al.* Prevalence of antineuronal antibodies in patients with encephalopathy of unknown etiology: data from a nationwide registry in Korea. *J Neuroimmunol* 2016; 293: 34–38.
  11. Lancaster E, Huijbers MG, Bar V, *et al.* Investigations of caspr2, an autoantigen of encephalitis and neuromyotonia. *Ann Neurol* 2011; 69: 303–311.
  12. Lai M, Hughes EG, Peng X, *et al.* AMPA receptor antibodies in limbic encephalitis alter synaptic receptor location. *Ann Neurol* 2009; 65: 424–434.
  13. Petit-Pedrol M, Armangue T, Peng X, *et al.* Encephalitis with refractory seizures, status epilepticus, and antibodies to the GABAA receptor: a case series, characterisation of the antigen, and analysis of the effects of antibodies. *Lancet Neurol* 2014; 13: 276–286.
  14. Lancaster E, Lai M, Peng X, *et al.* Antibodies to the GABA(B) receptor in limbic encephalitis with seizures: case series and characterisation of the antigen. *Lancet Neurol* 2010; 9: 67–76.
  15. Boronat A, Gelfand JM, Gresa-Arribas N, *et al.* Encephalitis and antibodies to dipeptidyl-peptidase-like protein-6, a subunit of Kv4.2 potassium channels. *Ann Neurol* 2013; 73: 120–128.
  16. Hutchinson M, Waters P, McHugh J, *et al.* Progressive encephalomyelitis, rigidity, and myoclonus: a novel glycine receptor antibody. *Neurology* 2008; 71: 1291–1292.
  17. Graus F, Titulaer MJ, Balu R, *et al.* A clinical approach to diagnosis of autoimmune encephalitis. *Lancet Neurol* 2016; 15: 391–404.
  18. Chevdeville A, Honnorat J, Hampe CS, *et al.* Neuronal central nervous system syndromes probably mediated by autoantibodies. *Eur J Neurosci* 2016; 43: 1535–1552.
  19. Malter MP, Helmstaedter C, Urbach H, *et al.* Antibodies to glutamic acid decarboxylase define a form of limbic encephalitis. *Ann Neurol* 2010; 67: 470–478.
  20. Arino H, Hoftberger R, Gresa-Arribas N, *et al.* Paraneoplastic neurological syndromes and glutamic acid decarboxylase antibodies. *JAMA Neurol* 2015; 72: 874–881.
  21. Fenalti G and Rowley MJ. GAD65 as a prototypic autoantigen. *J Autoimmun* 2008; 31: 228–232.
  22. Dalmau J, Geis C and Graus F. Autoantibodies to synaptic receptors and neuronal cell surface proteins in autoimmune diseases of the central nervous system. *Physiol Rev* 2017; 97: 839–887.
  23. Carvajal-Gonzalez A, Leite MI, Waters P, *et al.* Glycine receptor antibodies in PERM and related syndromes: characteristics, clinical features and outcomes. *Brain* 2014; 137: 2178–2192.
  24. Martinez-Hernandez E, Arino H, McKeon A, *et al.* Clinical and immunologic investigations in patients with stiff-person spectrum disorder. *JAMA Neurol* 2016; 73: 714–720.
  25. Sakaguchi S, Powrie F and Ransohoff RM. Re-establishing immunological self-tolerance in autoimmune disease. *Nat Med* 2012; 18: 54–58.
  26. Rawlings DJ, Metzler G, Wray-Dutra M, *et al.* Altered B cell signalling in autoimmunity. *Nat Rev Immunol* 2017; 17: 421–436.



27. Planaguma J, Leypoldt F, Mannara F, *et al.* Human N-methyl D-aspartate receptor antibodies alter memory and behaviour in mice. *Brain* 2015; 138: 94–109.
28. Wright S, Hashemi K, Stasiak L, *et al.* Epileptogenic effects of NMDAR antibodies in a passive transfer mouse model. *Brain* 2015; 138: 3159–3167.
29. Crisp SJ, Kullmann DM and Vincent A. Autoimmune synaptopathies. *Nat Rev Neurosci* 2016; 17: 103–117.
30. Hughes EG, Peng X, Gleichman AJ, *et al.* Cellular and synaptic mechanisms of anti-NMDA receptor encephalitis. *J Neurosci* 2010; 30: 5866–5875.
31. Moscato EH, Peng X, Jain A, *et al.* Acute mechanisms underlying antibody effects in anti-N-methyl-D-aspartate receptor encephalitis. *Ann Neurol* 2014; 76: 108–119.
32. Gleichman AJ, Spruce LA, Dalmau J, *et al.* Anti-NMDA receptor encephalitis antibody binding is dependent on amino acid identity of a small region within the GluN1 amino terminal domain. *J Neurosci* 2012; 32: 11082–11094.
33. Mikasova L, De Rossi P, Bouchet D, *et al.* Disrupted surface cross-talk between NMDA and Ephrin-B2 receptors in anti-NMDA encephalitis. *Brain* 2012; 135: 1606–1621.
34. Tuzun E, Zhou L, Baehring JM, *et al.* Evidence for antibody-mediated pathogenesis in anti-NMDAR encephalitis associated with ovarian teratoma. *Acta Neuropathol* 2009; 118: 737–743.
35. Martinez-Hernandez E, Horvath J, Shiloh-Malawsky Y, *et al.* Analysis of complement and plasma cells in the brain of patients with anti-NMDAR encephalitis. *Neurology* 2011; 77: 589–593.
36. Bien CG, Vincent A, Barnett MH, *et al.* Immunopathology of autoantibody-associated encephalitides: clues for pathogenesis. *Brain* 2012; 135: 1622–1638.
37. Irani SR, Pettingill P, Kleopa KA, *et al.* Morvan syndrome: clinical and serological observations in 29 cases. *Ann Neurol* 2012; 72: 241–255.
38. Arino H, Armangue T, Petit-Pedrol M, *et al.* Anti-LGI1-associated cognitive impairment: presentation and long-term outcome. *Neurology* 2016; 87: 759–765.
39. van Sonderen A, Arino H, Petit-Pedrol M, *et al.* The clinical spectrum of Caspr2 antibody-associated disease. *Neurology* 2016; 87: 521–528.
40. Kortvelyessy P, Bauer J, Stoppel CM, *et al.* Complement-associated neuronal loss in a patient with CASPR2 antibody-associated encephalitis. *Neurol Neuroimmunol Neuroinflamm* 2015; 2: e75.
41. Huijbers MG, Querol LA, Niks EH, *et al.* The expanding field of IgG4-mediated neurological autoimmune disorders. *Eur J Neurol* 2015; 22: 1151–1161.
42. Malter MP, Frisch C, Schoene-Bake JC, *et al.* Outcome of limbic encephalitis with VGKC-complex antibodies: relation to antigenic specificity. *J Neurol* 2014; 261: 1695–1705.
43. van Sonderen A, Thijs RD, Coenders EC, *et al.* Anti-LGI1 encephalitis: clinical syndrome and long-term follow-up. *Neurology* 2016; 87: 1449–1456.
44. Finke C, Pruss H, Heine J, *et al.* Evaluation of cognitive deficits and structural hippocampal damage in encephalitis with leucine-rich, glioma-inactivated 1 antibodies. *JAMA Neurol* 2017; 74: 50–59.
45. Miller TD, Chong TT, Aimola Davies AM, *et al.* Focal CA3 hippocampal subfield atrophy following LGI1 VGKC-complex antibody limbic encephalitis. *Brain*. Epub ahead of print 28 March 2017. DOI: 10.1093/brain/awx070.
46. Iizuka T, Yoshii S, Kan S, *et al.* Reversible brain atrophy in anti-NMDA receptor encephalitis: a long-term observational study. *J Neurol* 2010; 257: 1686–1691.
47. Iizuka T, Kaneko J, Tominaga N, *et al.* Association of progressive cerebellar atrophy with long-term outcome in patients with anti-N-methyl-D-aspartate receptor encephalitis. *JAMA Neurol* 2016; 73: 706–713.
48. Rother RP, Rollins SA, Mojcik CF, *et al.* Discovery and development of the complement inhibitor eculizumab for the treatment of paroxysmal nocturnal hemoglobinuria. *Nat Biotechnol* 2007; 25: 1256–1264.
49. Lee SK and Lee ST. The laboratory diagnosis of autoimmune encephalitis. *J Epilepsy Res* 2016; 6: 45–50.
50. van Coevorden-Hameete MH, Titulaer MJ, Schreurs MW, *et al.* Detection and characterization of autoantibodies to neuronal cell-surface antigens in the central nervous system. *Front Mol Neurosci* 2016; 9: 37.
51. Dalmau J, Tuzun E, Wu HY, *et al.* Paraneoplastic anti-N-methyl-D-aspartate receptor encephalitis associated with ovarian teratoma. *Ann Neurol* 2007; 61: 25–36.

52. Dalmau J, Gleichman AJ, Hughes EG, *et al.* Anti-NMDA-receptor encephalitis: case series and analysis of the effects of antibodies. *Lancet Neurol* 2008; 7: 1091–1098.
53. Lai M, Huijbers MG, Lancaster E, *et al.* Investigation of LGI1 as the antigen in limbic encephalitis previously attributed to potassium channels: a case series. *Lancet Neurol* 2010; 9: 776–785.
54. van Sonderen A, Schreurs MW, de Bruijn MA, *et al.* The relevance of VGKC positivity in the absence of LGI1 and Caspr2 antibodies. *Neurology* 2016; 86: 1692–1699.
55. Lang B, Makuch M, Moloney T, *et al.* Intracellular and non-neuronal targets of voltage-gated potassium channel complex antibodies. *J Neurol Neurosurg Psychiatry* 2017; 88: 353–361.
56. Gresa-Arribas N, Titulaer MJ, Torrents A, *et al.* Antibody titres at diagnosis and during follow-up of anti-NMDA receptor encephalitis: a retrospective study. *Lancet Neurol* 2014; 13: 167–177.
57. Wang R, Guan HZ, Ren HT, *et al.* CSF findings in patients with anti-N-methyl-D-aspartate receptor-encephalitis. *Seizure* 2015; 29: 137–142.
58. Kreye J, Wenke NK, Chayka M, *et al.* Human cerebrospinal fluid monoclonal N-methyl-D-aspartate receptor autoantibodies are sufficient for encephalitis pathogenesis. *Brain* 2016; 139: 2641–2652.
59. Rubenstein JL, Combs D, Rosenberg J, *et al.* Rituximab therapy for CNS lymphomas: targeting the leptomeningeal compartment. *Blood* 2003; 101: 466–468.
60. Hiepe F and Radbruch A. Plasma cells as an innovative target in autoimmune disease with renal manifestations. *Nat Rev Nephrol* 2016; 12: 232–240.
61. Behrendt V, Krogias C, Reinacher-Schick A, *et al.* Bortezomib treatment for patients with anti-N-methyl-d-aspartate receptor encephalitis. *JAMA Neurol* 2016; 73: 1251–1253.
62. Titulaer MJ, Kayser MS and Dalmau J. Authors' reply. *Lancet Neurol* 2013; 12: 425–426.
63. Scheibe F, Pruss H, Mengel AM, *et al.* Bortezomib for treatment of therapy-refractory anti-NMDA receptor encephalitis. *Neurology* 2017; 88: 366–370.
64. Iizuka T, Sakai F, Ide T, *et al.* Anti-NMDA receptor encephalitis in Japan: long-term outcome without tumor removal. *Neurology* 2008; 70: 504–511.
65. Szots M, Marton A, Kover F, *et al.* Natural course of LGI1 encephalitis: 3–5 years of follow-up without immunotherapy. *J Neurol Sci* 2014; 343: 198–202.
66. Dalmau J, Lancaster E, Martinez-Hernandez E, *et al.* Clinical experience and laboratory investigations in patients with anti-NMDAR encephalitis. *Lancet Neurol* 2011; 10: 63–74.
67. Hoftberger R, van Sonderen A, Leypoldt F, *et al.* Encephalitis and AMPA receptor antibodies: novel findings in a case series of 22 patients. *Neurology* 2015; 84: 2403–2412.
68. Finke C, Kopp UA, Pruss H, *et al.* Cognitive deficits following anti-NMDA receptor encephalitis. *J Neurol Neurosurg Psychiatry* 2012; 83: 195–198.
69. McKeon GL, Scott JG, Spooner DM, *et al.* Cognitive and social functioning deficits after anti-N-methyl-D-aspartate receptor encephalitis: an exploratory case series. *J Int Neuropsychol Soc* 2016; 22: 828–838.
70. Nosadini M, Mohammad SS, Ramanathan S, *et al.* Immune therapy in autoimmune encephalitis: a systematic review. *Expert Rev Neurother* 2015; 15: 1391–1419.
71. Irani SR, Bera K, Waters P, *et al.* N-methyl-D-aspartate antibody encephalitis: temporal progression of clinical and paraclinical observations in a predominantly non-paraneoplastic disorder of both sexes. *Brain* 2010; 133: 1655–1667.
72. Titulaer MJ, McCracken L, Gabilondo I, *et al.* Treatment and prognostic factors for long-term outcome in patients with anti-NMDA receptor encephalitis: an observational cohort study. *Lancet Neurol* 2013; 12: 157–165.
73. Kawasaki E. Type 1 diabetes and autoimmunity. *Clin Pediatr Endocrinol* 2014; 23: 99–105.
74. Barnes PJ. How corticosteroids control inflammation: Quintiles Prize Lecture 2005. *Br J Pharmacol* 2006; 148: 245–254.
75. Wiseman AC. Immunosuppressive medications. *Clin J Am Soc Nephrol* 2016; 11: 332–343.
76. Witt KA and Sandoval KE. Steroids and the blood-brain barrier: therapeutic implications. *Adv Pharmacol* 2014; 71: 361–390.
77. Slade JD and Hepburn B. Prednisone-induced alterations of circulating human lymphocyte subsets. *J Lab Clin Med* 1983; 101: 479–487.

78. Lahood N, Emerson SS, Kumar P, *et al.* Antibody levels and response to pneumococcal vaccine in steroid-dependent asthma. *Ann Allergy* 1993; 70: 289–294.
79. Lack G, Ochs HD and Gelfand EW. Humoral immunity in steroid-dependent children with asthma and hypogammaglobulinemia. *J Pediatr* 1996; 129: 898–903.
80. Vyas S, Rodrigues AJ, Silva JM, *et al.* Chronic stress and glucocorticoids: from neuronal plasticity to neurodegeneration. *Neural Plast* 2016; 2016: 6391686.
81. Jacob S and Rajabally YA. Current proposed mechanisms of action of intravenous immunoglobulins in inflammatory neuropathies. *Curr Neuropharmacol* 2009; 7: 337–342.
82. Hansen RJ and Balthasar JP. Intravenous immunoglobulin mediates an increase in anti-platelet antibody clearance via the FcRn receptor. *Thromb Haemost* 2002; 88: 898–899.
83. Li N, Zhao M, Hilario-Vargas J, *et al.* Complete FcRn dependence for intravenous Ig therapy in autoimmune skin blistering diseases. *J Clin Invest* 2005; 115: 3440–3450.
84. Gelfand EW. Intravenous immune globulin in autoimmune and inflammatory diseases. *N Engl J Med* 2012; 367: 2015–2025.
85. Iro MA, Sadarangani M, Absoud M, *et al.* Immunoglobulin in the treatment of encephalitis (IgNiTE): protocol for a multicentre randomised controlled trial. *BMJ Open* 2016; 6: e012356.
86. Reeves HM and Winters JL. The mechanisms of action of plasma exchange. *Br J Haematol* 2014; 164: 342–351.
87. DeSena AD, Noland DK, Matevosyan K, *et al.* Intravenous methylprednisolone versus therapeutic plasma exchange for treatment of anti-N-methyl-D-aspartate receptor antibody encephalitis: a retrospective review. *J Clin Apher* 2015; 30: 212–216.
88. Suppiej A, Nosadini M, Zuliani L, *et al.* Plasma exchange in pediatric anti-NMDAR encephalitis: a systematic review. *Brain Dev* 2016; 38: 613–622.
89. Klingel R, Heibges A and Fassbender C. Plasma exchange and immunoadsorption for autoimmune neurologic diseases - current guidelines and future perspectives. *Atheroscler Suppl* 2009; 10: 129–132.
90. Kohler W, Ehrlich S, Dohmen C, *et al.* Tryptophan immunoadsorption for the treatment of autoimmune encephalitis. *Eur J Neurol* 2015; 22: 203–206.
91. Dogan Onugoren M, Golombeck KS, Bien C, *et al.* Immunoadsorption therapy in autoimmune encephalitis. *Neurol Neuroimmunol Neuroinflamm* 2016; 3: e207.
92. Heine J, Ly LT, Lieker I, *et al.* Immunoadsorption or plasma exchange in the treatment of autoimmune encephalitis: a pilot study. *J Neurol* 2016; 263: 2395–2402.
93. Berger B, Hottenrott T, Leubner J, *et al.* Transient spurious intrathecal immunoglobulin synthesis in neurological patients after therapeutic apheresis. *BMC Neurol* 2015; 15: 255.
94. Lee WJ, Lee ST, Byun JI, *et al.* Rituximab treatment for autoimmune limbic encephalitis in an institutional cohort. *Neurology* 2016; 86: 1683–1691.
95. Lancaster E. The diagnosis and treatment of autoimmune encephalitis. *J Clin Neurol* 2016; 12: 1–13.
96. Kosmidis ML and Dalakas MC. Practical considerations on the use of rituximab in autoimmune neurological disorders. *Ther Adv Neurol Disord* 2010; 3: 93–105.
97. Kim SH, Kim W, Li XF, *et al.* Repeated treatment with rituximab based on the assessment of peripheral circulating memory B cells in patients with relapsing neuromyelitis optica over 2 years. *Arch Neurol* 2011; 68: 1412–1420.
98. Kim SH, Huh SY, Lee SJ, *et al.* A 5-year follow-up of rituximab treatment in patients with neuromyelitis optica spectrum disorder. *JAMA Neurol* 2013; 70: 1110–1117.
99. Damato V, Evoli A and Iorio R. Efficacy and safety of rituximab therapy in neuromyelitis optica spectrum disorders: a systematic review and meta-analysis. *JAMA Neurol* 2016; 73: 1342–1348.
100. Dalakas MC. Invited article: inhibition of B cell functions: implications for neurology. *Neurology* 2008; 70: 2252–2260.
101. Dale RC, Brilot F, Duffy LV, *et al.* Utility and safety of rituximab in pediatric autoimmune and inflammatory CNS disease. *Neurology* 2014; 83: 142–150.
102. Hachiya Y, Uruha A, Kasai-Yoshida E, *et al.* Rituximab ameliorates anti-N-methyl-D-aspartate receptor encephalitis by removal of short-lived plasmablasts. *J Neuroimmunol* 2013; 265: 128–130.

103. Brito-Zeron P, Kostov B, Bosch X, *et al.* Therapeutic approach to IgG4-related disease: a systematic review. *Medicine (Baltimore)* 2016; 95: e4002.
104. Sabater L, Gaig C, Gelpi E, *et al.* A novel non-rapid-eye movement and rapid-eye-movement parasomnia with sleep breathing disorder associated with antibodies to IgLON5: a case series, characterisation of the antigen, and post-mortem study. *Lancet Neurol* 2014; 13: 575–586.
105. Martin ST, Cardwell SM, Nailor MD, *et al.* Hepatitis B reactivation and rituximab: a new boxed warning and considerations for solid organ transplantation. *Am J Transplant* 2014; 14: 788–796.
106. Carson KR, Evens AM, Richey EA, *et al.* Progressive multifocal leukoencephalopathy after rituximab therapy in HIV-negative patients: a report of 57 cases from the Research on Adverse Drug Events and Reports project. *Blood* 2009; 113: 4834–4840.
107. Coiffier B, Lepage E, Briere J, *et al.* CHOP chemotherapy plus rituximab compared with CHOP alone in elderly patients with diffuse large-B-cell lymphoma. *N Engl J Med* 2002; 346: 235–242.
108. Gourley MF, Austin HA III, Scott D, *et al.* Methylprednisolone and cyclophosphamide, alone or in combination, in patients with lupus nephritis. A randomized, controlled trial. *Ann Intern Med* 1996; 125: 549–557.
109. Jayne D, Rasmussen N, Andrassy K, *et al.* A randomized trial of maintenance therapy for vasculitis associated with antineutrophil cytoplasmic autoantibodies. *N Engl J Med* 2003; 349: 36–44.
110. Hickman LC, Valentine LN and Falcone T. Preservation of gonadal function in women undergoing chemotherapy: a review of the potential role for gonadotropin-releasing hormone agonists. *Am J Obstet Gynecol* 2016; 215: 415–422.
111. Ginzler EM and Aranow C. Mycophenolate mofetil in lupus nephritis. *Lupus* 2005; 14: 59–64.
112. Lee WJ, Lee ST, Moon J, *et al.* Tocilizumab in autoimmune encephalitis refractory to rituximab: An Institutional Cohort Study. *Neurotherapeutics* 2016; 13: 824–832.
113. Lim JA, Lee ST, Moon J, *et al.* New feasible treatment for refractory autoimmune encephalitis: low-dose interleukin-2. *J Neuroimmunol* 2016; 299: 107–111.
114. Kang S, Tanaka T and Kishimoto T. Therapeutic uses of anti-interleukin-6 receptor antibody. *Int Immunol* 2015; 27: 21–29.
115. Mihara M, Kasutani K, Okazaki M, *et al.* Tocilizumab inhibits signal transduction mediated by both mIL-6R and sIL-6R, but not by the receptors of other members of IL-6 cytokine family. *Int Immunopharmacol* 2005; 5: 1731–1740.
116. Smolen JS, Beaulieu A, Rubbert-Roth A, *et al.* Effect of interleukin-6 receptor inhibition with tocilizumab in patients with rheumatoid arthritis (OPTION study): a double-blind, placebo-controlled, randomised trial. *Lancet* 2008; 371: 987–997.
117. De Benedetti F, Brunner HI, Ruperto N, *et al.* Randomized trial of tocilizumab in systemic juvenile idiopathic arthritis. *N Engl J Med* 2012; 367: 2385–2395.
118. Ayzenberg I, Kleiter I, Schroder A, *et al.* Interleukin 6 receptor blockade in patients with neuromyelitis optica nonresponsive to anti-CD20 therapy. *JAMA Neurol* 2013; 70: 394–397.
119. Araki M, Matsuoka T, Miyamoto K, *et al.* Efficacy of the anti-IL-6 receptor antibody tocilizumab in neuromyelitis optica: a pilot study. *Neurology* 2014; 82: 1302–1306.
120. Ringelstein M, Ayzenberg I, Harmel J, *et al.* Long-term therapy with interleukin 6 receptor blockade in highly active neuromyelitis optica spectrum disorder. *JAMA Neurol* 2015; 72: 756–763.
121. Krogias C, Hoepner R, Muller A, *et al.* Successful treatment of anti-Caspr2 syndrome by interleukin 6 receptor blockade through tocilizumab. *JAMA Neurol* 2013; 70: 1056–1059.
122. Shetty A, Hanson R, Korsten P, *et al.* Tocilizumab in the treatment of rheumatoid arthritis and beyond. *Drug Des Devel Ther* 2014; 8: 349–364.
123. Rossi JF, Lu ZY, Jourdan M, *et al.* Interleukin-6 as a therapeutic target. *Clin Cancer Res* 2015; 21: 1248–1257.
124. Buckner JH. Mechanisms of impaired regulation by CD4(+)CD25(+)FOXP3(+) regulatory T cells in human autoimmune diseases. *Nat Rev Immunol* 2010; 10: 849–859.
125. Carbone F, De Rosa V, Carrieri PB, *et al.* Regulatory T cell proliferative potential is impaired in human autoimmune disease. *Nat Med* 2014; 20: 69–74.




126. Malek TR and Castro I. Interleukin-2 receptor signaling: at the interface between tolerance and immunity. *Immunity* 2010; 33: 153–165.
127. Liao W, Lin JX and Leonard WJ. Interleukin-2 at the crossroads of effector responses, tolerance, and immunotherapy. *Immunity* 2013; 38: 13–25.
128. Yu A, Zhu L, Altman NH, *et al.* A low interleukin-2 receptor signaling threshold supports the development and homeostasis of T regulatory cells. *Immunity* 2009; 30: 204–217.
129. Yu A, Snowwhite I, Vendrame F, *et al.* Selective IL-2 responsiveness of regulatory T cells through multiple intrinsic mechanisms supports the use of low-dose IL-2 therapy in type 1 diabetes. *Diabetes* 2015; 64: 2172–2183.
130. Klatzmann D and Abbas AK. The promise of low-dose interleukin-2 therapy for autoimmune and inflammatory diseases. *Nat Rev Immunol* 2015; 15: 283–294.
131. Saadoun D, Rosenzweig M, Joly F, *et al.* Regulatory T-cell responses to low-dose interleukin-2 in HCV-induced vasculitis. *N Engl J Med* 2011; 365: 2067–2077.
132. Castela E, Le Duff F, Butori C, *et al.* Effects of low-dose recombinant interleukin 2 to promote T-regulatory cells in alopecia areata. *JAMA Dermatol* 2014; 150: 748–751.
133. He J, Zhang X, Wei Y, *et al.* Low-dose interleukin-2 treatment selectively modulates CD4(+) T cell subsets in patients with systemic lupus erythematosus. *Nat Med* 2016; 22: 991–993.
134. Koreth J, Matsuoka K, Kim HT, *et al.* Interleukin-2 and regulatory T cells in graft-versus-host disease. *N Engl J Med* 2011; 365: 2055–2066.
135. Meister S, Schubert U, Neubert K, *et al.* Extensive immunoglobulin production sensitizes myeloma cells for proteasome inhibition. *Cancer Res* 2007; 67: 1783–1792.
136. Gabilondo I, Saiz A, Galan L, *et al.* Analysis of relapses in anti-NMDAR encephalitis. *Neurology* 2011; 77: 996–999.
137. Uzawa A, Mori M, Takahashi Y, *et al.* Anti-N-methyl D-aspartate-type glutamate receptor antibody-positive limbic encephalitis in a patient with multiple sclerosis. *Clin Neurol Neurosurg* 2012; 114: 402–404.
138. Takeda A, Shimada H, Tamura A, *et al.* A case of anti-N-methyl-d-aspartate receptor encephalitis with multiple sclerosis-like demyelinated lesions. *Mult Scler Relat Disord* 2014; 3: 391–397.
139. Fleischmann R, Pruss H, Rosche B, *et al.* Severe cognitive impairment associated with intrathecal antibodies to the NR1 subunit of the N-methyl-D-aspartate receptor in a patient with multiple sclerosis. *JAMA Neurol* 2015; 72: 96–99.
140. Kruer MC, Koch TK, Bourdette DN, *et al.* NMDA receptor encephalitis mimicking seronegative neuromyelitis optica. *Neurology* 2010; 74: 1473–1475.
141. Titulaer MJ, Hofberger R, Iizuka T, *et al.* Overlapping demyelinating syndromes and anti-N-methyl-D-aspartate receptor encephalitis. *Ann Neurol* 2014; 75: 411–428.
142. Gotterer L and Li Y. Maintenance immunosuppression in myasthenia gravis. *J Neurol Sci* 2016; 369: 294–302.
143. Papadopoulos MC, Bennett JL and Verkman AS. Treatment of neuromyelitis optica: state-of-the-art and emerging therapies. *Nat Rev Neurol* 2014; 10: 493–506.
144. Steinman L, Bar-Or A, Behne JM, *et al.* Restoring immune tolerance in neuromyelitis optica: Part I. *Neurol Neuroimmunol Neuroinflamm* 2016; 3: e276.
145. Bar-Or A, Steinman L, Behne JM, *et al.* Restoring immune tolerance in neuromyelitis optica: Part II. *Neurol Neuroimmunol Neuroinflamm* 2016; 3: e277.
146. de Montmollin E, Demeret S, Brule N, *et al.* Anti-N-methyl-D-aspartate receptor encephalitis in adult patients requiring intensive care. *Am J Respir Crit Care Med* 2017; 195: 491–499.
147. Byun JI, Lee ST, Jung KH, *et al.* Effect of immunotherapy on seizure outcome in patients with autoimmune encephalitis: a prospective observational registry study. *PLoS One* 2016; 11: e0146455.
148. Hansen HC, Klingbeil C, Dalmau J, *et al.* Persistent intrathecal antibody synthesis 15 years after recovering from anti-N-methyl-D-aspartate receptor encephalitis. *JAMA Neurol* 2013; 70: 117–119.
149. Kowarik MC, Cepok S, Sellner J, *et al.* CXCL13 is the major determinant for B cell recruitment to the CSF during neuroinflammation. *J Neuroinflammation* 2012; 9: 93.
150. Leyboldt F, Hofberger R, Titulaer MJ, *et al.* Investigations on CXCL13 in anti-N-methyl-D-aspartate receptor encephalitis: a potential



- biomarker of treatment response. *JAMA Neurol* 2015; 72: 180–186.
151. Kothur K, Wienholt L, Mohammad SS, *et al.* Utility of CSF cytokine/chemokines as markers of active intrathecal inflammation: comparison of demyelinating, anti-NMDAR and enteroviral encephalitis. *PLoS One* 2016; 11: e0161656.
152. Constantinescu R, Krysl D, Bergquist F, *et al.* Cerebrospinal fluid markers of neuronal and glial cell damage to monitor disease activity and predict long-term outcome in patients with autoimmune encephalitis. *Eur J Neurol* 2016; 23: 796–806.
153. Pettingill P, Kramer HB, Coebergh JA, *et al.* Antibodies to GABAA receptor alpha1 and gamma2 subunits: clinical and serologic characterization. *Neurology* 2015; 84: 1233–1241.
154. Hoftberger R, Titulaer MJ, Sabater L, *et al.* Encephalitis and GABAB receptor antibodies: novel findings in a new case series of 20 patients. *Neurology* 2013; 81: 1500–1506.
155. Tobin WO, Lennon VA, Komorowski L, *et al.* DPPX potassium channel antibody: frequency, clinical accompaniments, and outcomes in 20 patients. *Neurology* 2014; 83: 1797–1803.

Visit SAGE journals online  
[journals.sagepub.com/  
home/tan](http://journals.sagepub.com/home/tan)

 SAGE journals